

Original Article

The peptide glycyl-L-histidyl-L-lysine is an endogenous antioxidant in living organisms, possibly by diminishing hydroxyl and peroxy radicals

Satoru Sakuma, Mai Ishimura, Yukinori Yuba, Yuhki Itoh, Yohko Fujimoto

Department of Physiological Chemistry, Osaka University of Pharmaceutical Sciences, 4-20-1 Nasahara, Takatsuki, Osaka 569-1094, Japan

Received May 13, 2018; Accepted June 16, 2018; Epub June 25, 2018; Published June 30, 2018

Abstract: Despite evidence that tripeptide glycyl-L-histidyl-L-lysine (GHK) is an endogenous antioxidant, its mechanism and importance are not fully understood. In the present study, the ability of GHK to reduce levels of reactive oxygen species (ROS) in Caco-2 cells was evaluated by flow cytometry with the oxidation-sensitive fluorescent dye 2',7'-dichlorodihydrofluorescein diacetate. Further, types of ROS diminished by GHK were assessed by utilizing an electron spin resonance (ESR) spin-trapping technique. GHK reduced the *tert*-butyl hydroperoxide-induced increase in ROS levels in Caco-2 cells at concentrations of 10 μ M or less. Experiments utilizing an ESR spin-trapping technique revealed that, among hydroxyl (\cdot OH), superoxide ($O_2^{\cdot-}$), and peroxy (ROO \cdot) radicals generated by respective chemical reaction systems, GHK diminished signals of both \cdot OH and ROO \cdot , but not $O_2^{\cdot-}$. Additionally, the GHK effect on the signal of \cdot OH was much stronger than those of other well-known antioxidative, endogenous peptides, carnosine and reduced glutathione. These results suggest that GHK can function as an endogenous antioxidant in living organisms, possibly by diminishing \cdot OH and ROO \cdot .

Keywords: Glycyl-L-histidyl-L-lysine, hydroxyl radical, peroxy radical, oxidative stress, endogenous antioxidant

Introduction

The natural tripeptide glycyl-L-histidyl-L-lysine (GHK) was first isolated by Pickart and Thaler [1]. GHK is liberated from extracellular matrix proteins, especially the α -II chain of collagen in response to soft tissue damage [2-4]. Numerous studies have demonstrated that this simple molecule improves wound healing and tissue regeneration (e.g., skin, hair follicles, bones, stomach, intestinal linings, and liver tissues), increases collagen and glycosaminoglycans, and increases angiogenesis and nerve outgrowth [5-8]. Interestingly, Bobyntsev et al. [9] also reported that the administration of GHK has marked anxiolytic effects on behavioral responses of rats. Thus, GHK has many beneficial effects on the body, but the mechanisms underlying these effects are not fully understood.

Substantial evidence has surfaced regarding the harmful effects of reactive oxygen species

(ROS) generated endogenously and exogenously. Certain antioxidants may have physiological and pharmacological functions, without any side effects, in preventing oxidative stress-induced damage. GHK has been reported to quench the toxic products of lipid peroxidation, α,β -4-hydroxy-trans-2-nonenal and acrolein, which play important roles in the pathogenesis of several age-related conditions [10, 11]. However, to the best of our knowledge, the effect of GHK on ROS levels in living cells and on various types of ROS are unknown.

In this study, we examined the concentration-dependent effects of GHK on ROS levels in Caco-2 cells by using flow cytometry with the oxidation-sensitive fluorescent dye 2',7'-dichlorodihydrofluorescein diacetate (DCFH-DA). We also utilized an electron-spin resonance spectrometry (ESR) spin-trapping technique [12] to evaluate the specific effects of GHK on hydroxyl (\cdot OH), superoxide ($O_2^{\cdot-}$), and peroxy (ROO \cdot) radicals generated by respective chemical reaction

Diminution of intracellular ROS level by GHK

systems and compared these effects with those of other well-known antioxidative, endogenous peptides, carnosine and reduced glutathione (GSH) [13, 14].

Materials and methods

Materials

GHK, H₂O₂, mannitol, SOD, 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox), xanthine, xanthine oxidase (XO), *tert*-butyl hydroperoxide (t-BOOH, 70%, w/w, aqueous solution), GSH, and diethylenetriaminepentaacetic acid (DTPA) were obtained from Sigma Chemical Co. (St. Louis, MO, USA). DCFH-DA was obtained from Life Technologies Corporation (Carlsbad, CA, USA). 5,5-Dimethyl-1-pyrroline-*N*-oxide (DMPO) and α -(4-pyridyl 1-oxide)-*N*-*tert*-butylnitron (POBN) were purchased from Enzo Life Technologies (Farmingdale, NY, USA). *N*-Acetyl-L-cysteine (NAC), Ce(SO₄)₂·4H₂O, and FeSO₄·7H₂O were obtained from Wako Pure Chemical Industries, Limited (Osaka, Japan). All other reagents were of analytical grade. The ultrapure water was prepared using a compact ultrapure water system (Milli-Q[®]; Merck Millipore, Burlington, MA, USA).

Measurement of ROS in Caco-2 cells

Measurements of ROS in Caco-2 cells were performed following previously reported methods [15]. Caco-2 cells were purchased from the European Collection of Cell Cultures (Salisbury, Wilts, UK) and cultured in Minimum Essential Medium (Life Technologies Corporation) supplemented with 10% fetal bovine serum (Nichirei Biosciences Inc., Tokyo, Japan) and 1% non-essential amino acids (Life Technologies Corporation). The cells were maintained in a humidified atmosphere of 5% carbon dioxide at 37°C.

ROS levels were measured by flow cytometry with the oxidation-sensitive fluorescent dye, DCFH-DA. The cells (1.0 × 10⁶ cells/28 cm² dish) were incubated with the test reagents for 6 h, and then DCFH-DA was added at a final concentration of 10 μM. After incubation for 30 min, the cells were collected by centrifugation (4°C and 200 × *g* for 5 min) and washed twice. The samples were filtered through a nylon mesh (37 μm) and subjected to flow cytometry (FACS Aria™ III Flow Cytometer; Becton Dickinson, Basel, Switzerland).

ESR measurement

The ESR spectrometer TE-2100 (JEOL, Tokyo, Japan) and a JEOL flat quartz cell were used. The conditions were as follows: field, 336 ± 5 mT width; power, 2 mW (DMPO-OH· signal) and 4 mW (DMPO-OOH·, and POBN adduct signals); field modulation, 0.200 mT; time constant, 0.1; and amplitude, 300. A manganese signal was used as the external standard.

Observation of DMPO-OH· reflecting ·OH

The Fenton reaction was initiated by adding H₂O₂ (final concentration, 0.5 mM) to a mixture of DMPO (final concentration, 10 mM) and FeSO₄·7H₂O (final concentration, 0.25 mM) in 0.1 M sodium phosphate buffer (pH 7.4) in a total volume of 0.5 mL. The spin-trapped DMPO-OH· signal reflecting ·OH was measured 1 min after the addition of H₂O₂.

Observation of DMPO-OOH· reflecting O₂·-

The xanthine/XO reaction was started by adding xanthine (final concentration, 200 mM) to a mixture of DMPO (final concentration, 220 mM), XO (final concentration, 0.1 U/mL), and DTPA (final concentration, 0.1 mM) in 0.1 M sodium phosphate buffer (pH 7.4) in a total volume of 0.5 mL. The spin-trapped DMPO-OOH· signal reflecting O₂·- was measured 1 min after the addition of xanthine.

Observation of POBN-adduct signal reflecting t-BOO·

The Ce⁴⁺/t-BOOH reaction was started by adding t-BOOH (final concentration, 400 mM) to a mixture of POBN (final concentration, 10 mM) and Ce(SO₄)₂·4H₂O (final concentration, 0.2 mM) in 0.1 M sodium phosphate buffer (pH 7.4) in a total volume of 0.5 mL. The POBN adduct signal reflecting t-BOO· were measured 1 min after the addition of t-BOOH.

Statistical analysis

The results are expressed as means ± standard errors of the mean. Significant differences between two groups were assessed using *t*-tests, and differences between multiple groups were assessed by one-way analysis of variance, followed by Scheffé's multiple comparison tests. *P*-values less than 0.05 were considered statistically significant.

Diminution of intracellular ROS level by GHK




Figure 1. Alterations in ROS levels in Caco-2 cells treated with t-BOOH in combination with GHK. (A) Representative univariate histograms of flow cytometry data. (B) Quantitative assessment of the MFI flow cytometry data. Results are presented as means \pm standard errors of the mean ($n = 3-12$). $^{\alpha}P < 0.01$ vs. None. $^{\beta}P < 0.01$, $^{\gamma}P < 0.01$ vs. t-BOOH. $^{\delta}P < 0.01$ vs. NAC. MFI, mean fluorescence intensity (shown as dashed lines in part A); GHK, glycyl-L-histidyl-L-lysine; t-BOOH, tert-butylhydroperoxide; NAC, N-acetyl-L-cysteine.

Results and discussion

GHK potently diminishes ROS levels in viable cells

Figure 1A shows the effects of t-BOOH with or without GHK and NAC on intracellular ROS generation in Caco-2 cells, as determined by flow cytometry with the redox-sensitive fluorescent dye DCFH-DA. NAC is often used as an antioxidant in cell experiments [16]. The addition of t-BOOH (50 μ M) to Caco-2 cells shifted the mean fluorescence intensity (MFI, dashed lines in **Figure 1A**) to the right, indicating an increase in ROS levels based on DCF fluorescence. The increment in MFI induced by t-BOOH was reduced by the addition of 10 μ M GHK. **Figure 1B** summarizes the MFI results obtained by the method described in **Figure 1A**. The addition of 10 μ M NAC significantly diminished the t-BOOH-induced increase in MFI (28% inhibition). At the same concentration, GHK significantly reduced

the t-BOOH-induced increase in MFI (10 μ M GHK, 48% inhibition). The inhibitory effect of 10 μ M GHK was stronger than that of NAC, with a statistically significance. This means that GHK reduces the t-BOOH-induced increase in ROS levels in living cells, even at concentrations of 10 μ M or less.

GHK diminishes the amount of spin signal adduct of \cdot OH more potently than do carnosine and GSH

A direct method for measuring free radicals in aqueous conditions is ESR spectroscopy [14]. **Figure 2A** shows the ESR spectra of the spin signal adducts from \cdot OH, $O_2^{\cdot-}$, and t-BOO \cdot using their respective spin trapping reagents.

Figure 2Aa shows the ESR spectra obtained by the Fenton reaction and DMPO with and without mannitol, a specific \cdot OH scavenger [17, 18]. The 1:2:2:1 quartet pattern shows a hyperfine

Diminution of intracellular ROS level by GHK




Figure 2. Effect of GHK on the amounts of spin signal adducts of $\cdot\text{OH}$, $\text{O}_2\cdot^-$, and t-BOO \cdot generated by their respective chemical reaction systems. A. Representative ESR signal spectra of (a) DMPO-OH reflecting $\cdot\text{OH}$, (b) DMPO-OOH reflecting $\text{O}_2\cdot^-$, and (c) POBN adduct signal reflecting t-BOO \cdot . (a) Fe^{2+} , 0.25 mM; H_2O_2 , 0.5 mM. (b) xanthine, 200 mM; XO, 0.1 U/mL. (c) Ce^{4+} , 0.2 mM; t-BOOH, 400 mM. B. The effect of GHK on the amounts of spin signal adducts of $\cdot\text{OH}$, $\text{O}_2\cdot^-$, and t-BOO \cdot . The radical intensity was defined as the ratio of the peak height of respective signal [indicated as arrows in part Aa-Ac] to that of Manganese (Mn). Results are presented as means \pm standard errors of the mean ($n = 3-8$). ^α $P < 0.01$ vs. Control. GHK, 250 μM . $\cdot\text{OH}$, hydroxyl radicals; $\text{O}_2\cdot^-$, superoxide radicals; t-BOO \cdot , *tert*-butyl peroxy radicals; GHK, glycyl-L-histidyl-L-lysine; XO, xanthine oxidase; SOD, superoxide dismutase; ESR, electron spin resonance; t-BOOH, *tert*-butyl hydroperoxide.

coupling at 1.50 mT, the same value as that of DMPO-OH \cdot reported previously [19, 20]. The signal of DMPO-OH \cdot was quenched almost completely with 10 mM mannitol. **Figure 2Ab** shows the spin signal adducts of DMPO-OOH \cdot reflecting $\text{O}_2\cdot^-$ in the absence or presence of 0.5 U/mL SOD. The hyperfine fit parameters for the DMPO adducts were as follows: $a(\text{N}) = 1.42$ mT, $a(\text{H})^\beta = 1.14$ mT, and $a(\text{H})^\gamma = 0.12$ mT, consistent with previous values reported for DMPO-OOH \cdot [19, 20]. The involvement of $\text{O}_2\cdot^-$ was confirmed by the reduction in the DMPO-OOH \cdot signal intensity in the presence of 0.5 U/mL SOD. **Figure 2Ac** shows the spin signal adduct of t-BOO \cdot [$a(\text{N}) = 1.51$ mT, $a(\text{H}) = 0.23$ mT] by the reaction of the Ce^{4+} /t-BOOH system with POBN. The hyperfine fit parameters are identical to those previously reported [21, 22]. An obvious quenching by 0.5 mM Trolox, which is a potent peroxy radical scavenger [23, 24], supported the identity of the product. Positive correlations between the disappearance of the signals of DMPO-OH \cdot , DMPO-OOH \cdot , and POBN adduct signal of t-BOO \cdot , and their respective scavenger concentrations were observed ([Supplementary Data](#)). **Figure 2A** and [Supplementary Data](#) collectively dem-

onstrate that these three experimental conditions using the ESR apparatus can be used to assess the quenching efficacies of specific substances against $\cdot\text{OH}$, $\text{O}_2\cdot^-$, and t-BOO \cdot . **Figure 2A** also showed representative ESR spectra in the presence of GHK. GHK at 250 μM reduced both spin signal adducts of $\cdot\text{OH}$ (1) and t-BOO \cdot (3), but not of $\text{O}_2\cdot^-$ (2).

Figure 2B shows the effect of GHK on the amounts of spin signal adducts of $\cdot\text{OH}$ (1), $\text{O}_2\cdot^-$ (2), and t-BOO \cdot (3) detected by the methods described in **Figure 2Aa-Ac**. GHK at 250 μM quenched the spin signal adducts of both $\cdot\text{OH}$ (47% inhibition) and t-BOO \cdot (30% inhibition), but did not influence the spin signal adduct of $\text{O}_2\cdot^-$.

The effect of GHK on the amount of spin signal adduct of $\cdot\text{OH}$ was investigated further. The diminishing effect of GHK on the amount of spin signal adduct of $\cdot\text{OH}$ was compared to that of an equimolar mixture (G+H+K, 1:1:1) of the GHK constitutive amino acids (**Table 1**). A mixture of G, H, and K (250 μM each) did not have any effect on the spin signal adduct of $\cdot\text{OH}$, whereas GHK at the same concentration did.

Diminution of intracellular ROS level by GHK




Figure 3. Effects of carnosine and GSH as well as GHK on the amounts of spin signal adduct of $\cdot\text{OH}$ generated by the Fenton-type reaction. The radical intensity was defined as the ratio of the peak height of the signal [indicated as arrows in **Figure 2Aa**] to that of Manganese (Mn). Results are presented as means \pm standard errors of the mean ($n = 3-8$). ^α $P < 0.05$, ^β $P < 0.01$ vs. None., ^γ $P < 0.01$ vs. GSH. ^δ $P < 0.05$, ^ϵ $P < 0.01$ vs. carnosine. $\cdot\text{OH}$, hydroxyl radicals; GHK, glycyl-L-histidyl-L-lysine; GSH, reduced glutathione.

Table 1. Effects of G plus H plus K as well as GHK on the generation of hydroxyl radicals

Treatment	OH (% of control)
Control	100.0 \pm 3.6
GHK	71.4 \pm 4.4 ^{α,β}
G+H+K	104.6 \pm 5.0

The radical intensity was defined as the ratio of the peak height of the signal [indicated as arrows in **Figure 2Aa-Ac**] to that of Manganese (Mn). Results are presented as means \pm standard errors of the mean ($n = 4-10$). ^α $P < 0.01$, significantly different from Control, ^β $P < 0.01$; significantly different from G+H+K. GHK, 250 μM ; G, H or K, 250 μM . G, glycine; H, histidine; K, lysine.

These results indicate that GHK results in a cooperative or synergistic reduction in the spin signal adduct of $\cdot\text{OH}$ when the amino acids were linked in the peptide structure.

Many studies have examined the antioxidant and metal ion chelation effects of carnosine (β -alanyl-L-histidine), a dipeptide, and GSH (γ -glutamyl-cysteinyl-glycine), a tri-peptide [13, 14]. **Figure 3** illustrates the $\cdot\text{OH}$ diminishing efficacy of GHK in comparison with those of carnosine and GSH. All three small peptides dose-dependently diminished $\cdot\text{OH}$, but the effect of GHK was much stronger than those of the other two peptides (IC_{50} value: GHK, approximately

250 μM ; carnosine, approximately 500 μM ; GSH, greater than 1000 μM). This finding shows that GHK diminishes the spin signal adduct of $\cdot\text{OH}$ more strongly than do carnosine and GSH.

GHK has been co-isolated from human plasma in association with the albumin and α -globulin fractions at about 200 ng/mL (0.6 μM) [1]. However, GHK has been reported to be liberated from extracellular matrix proteins, especially the α -II chain of collagen in response to even soft tissue damage [2-4]; accordingly, GHK level is likely to be detected at a broad range of concentrations, from nanomolar up to micromolar under certain pathophysiological conditions.

The present study showed that GHK, even at concentration of 10 μM or less, significantly reduces ROS levels initiated by t-BOOH. Thus, it is possible that GHK acts as an antioxidant under such pathophysiological conditions.

Acrolein, a well-known carbonyl toxin, is produced by lipid peroxidation of polyunsaturated fatty acids. GHK has been reported to effectively reduce the formation of both acrolein and another product of oxidation, 4-hydroxynonenal [10, 11]. These previous studies make us expect a direct scavenging capacity of GHK against $\cdot\text{OH}$ and $\text{ROO}\cdot$. But, it has been shown that GHK interacts with an approximately equimolar amount of copper and an approximately 1/5 molar amount of iron [17, 25]. Accordingly, it seems likely that GHK chelates Fe and Cu in the chemical reaction systems for the generation of $\cdot\text{OH}$ and $\text{ROO}\cdot$, and this effect is stronger than those of carnosine and GSH. Additional studies are needed to clarify the mechanism by which GHK diminishes the generation of $\cdot\text{OH}$ and $\text{ROO}\cdot$.

Conclusion

GHK significantly reduced t-BOOH-induced increases in ROS levels in Caco-2 cells, and this effect was stronger than that of NAC. GHK also diminished the generation of $\cdot\text{OH}$ and $\text{ROO}\cdot$ by

Diminution of intracellular ROS level by GHK

using an ESR method. We believe that our study makes a significant contribution to the literature because, to the best of our knowledge, this is the first time that GHK has the potential to reduce oxidative stress in living organisms, possibly by diminishing ·OH and ROO·.

Disclosure of conflict of interest

None.

Address correspondence to: Yohko Fujimoto, Department of Physiological Chemistry, Osaka University of Pharmaceutical Sciences, 4-20-1 Nasahara, Takatsuki, Osaka 569-1094, Japan. E-mail: fujimoto@gly.oups.ac.jp

References

- [1] Pickart L, Thaler MM. Tripeptide in human serum which prolongs survival of normal liver cells and stimulates growth in neoplastic liver. *Nat New Biol* 1973; 234: 85-87.
- [2] Pickart L, Vasquez-Soltero JM, Margolina A. The human tripeptide GHK-Cu in prevention of oxidative stress and degenerative conditions of aging: implications for cognitive health. *Oxid Med Cell Longev* 2012; 2012: 324832-324839.
- [3] Kukowska M, Dzierzbicka K. Developments in the synthesis and biological activity of Glycyl-L-Histidyl-L-Lysine derivatives. *Curr Med Chem* 2014; 221: 1505-1521.
- [4] Kukowska M, Kukowska-Kaszuba M, Dzierzbicka K. In vitro studies of antimicrobial activity of Gly-His-Lys conjugates as potential and promising candidates for therapeutics in skin and tissue infections. *Bioorg Med Chem Lett* 2015; 25: 542-546.
- [5] Pickart L. The human tri-peptide GHK and tissue remodeling. *J Biomater Sci Polym Ed* 2008; 19: 969-988.
- [6] Pickart L. The human tripeptide GHK (Glycyl-L-Histidyl-L-Lysine). The copper switch, and the treatment of the degenerative conditions of aging. In: Klatz R, Goldman R, editors. *American Academy of Anti-Aging Medicine* 2009. pp. 301-312.
- [7] Choi HR, Kang YA, Ryoo SJ, Shin JW, Na JI, Huh CH, Park KC. Stem cell recovering effect of copper-free GHK in skin. *J Pept Sci* 2012; 18: 685-690.
- [8] Jose S, Hughbanks ML, Binder BY, Ingavle GC, Leach JK. Enhanced trophic factor secretion by mesenchymal stem/stromal cells with Glycine-Histidine-Lysine (GHK)-modified alginate hydrogels. *Acta Biomater* 2014; 10: 1955-1964.
- [9] Bobyntsev II, Chernysheva OI, Dolgintsev ME, Smakhtin MY, Belykh AE. Anxiolytic effects of Gly-His-Lys peptide and its analogs. *Bull Exp Biol Med* 2015; 158: 726-728.
- [10] Beretta G, Artali R, Regazzoni L, Panigati M, Facino RM. Glycyl-histidyl-lysine (GHK) is a quencher of α,β -4-Hydroxy-*trans*-2-nonenal: a comparison with carnosine. Insights into the mechanism of reaction by electrospray ionization mass spectrometry, ^1H NMR, and computational techniques. *Chem Res Toxicol* 2007; 20: 1309-1314.
- [11] Beretta G, Arlandini E, Artali R, Anton JM, Maffei Facino R. Acrolein sequestering ability of the endogenous tripeptide glycyl-histidyl-lysine (GHK): characterization of conjugation products by ESI-MSⁿ and theoretical calculations. *J Pharm Biomed Anal* 2008; 47: 596-602.
- [12] Venkataraman S, Schafer FQ, Buettner GR. Detection of lipid radicals using EPR. *Antioxid Redox Singal* 2004; 6: 631-638.
- [13] Decker EA, Livisay SA, Zhou S. A re-evaluation of the antioxidant activity of purified carnosine. *Biochemistry (Moscow)* 2000; 65: 901-906.
- [14] Lluís JM, Morales A, Blasco C, Colell A, Mari M, Garcia-Ruiz C, Fernandez-Checa JC. Critical role of mitochondrial glutathione in the survival of hepatocytes during hypoxia. *J Biol Chem* 2005; 280: 3224-3232.
- [15] Kohda T, Sakuma S, Fujimoto Y. Vitamin E-like molecules potentiate the curcumin-induced suppression of Caco-2 cell proliferation. *Int J Pharmacol Res* 2016; 6: 41-46.
- [16] Lasram MM, Dhoub IB, Annabi A, El Fazaa S, Gharbi N. A review on the possible molecular mechanism of action of N-acetylcysteine against insulin resistance and type-2 diabetes development. *Clin Biochem* 2015; 48: 1200-1208.
- [17] Duffy SJ, Biegelsen ES, Holbrook M, Russell JD, Gokce N, Keaney JF Jr, Vita JA. Iron chelation improves endothelial function in patients with coronary artery disease. *Circulation* 2001; 103: 2799-2804.
- [18] Nakagawa S. Copper uptake is required for pyrrolidine dithiocarbamate-mediated oxidation and protein level increase of p53 in cells. Estimation of relative reaction rate of hydroxy radical with poly-hydroxy benzenes: ESR spin trapping combined with UV-A photolysis. *Anal Sci* 2013; 29: 377-380.
- [19] Buettner GR. The spin trapping of superoxide and hydroxyl free radicals with DMPO (5,5-Dimethylpyrrolidine-N-oxide): more about iron. *Free Rad Res Commun* 1993; 19: S79-S87.
- [20] Rangelova K, Mason RP. The fidelity of spin trapping with DMPO in biological systems. *Magn Reson Chem* 2011; 49: 152-158.
- [21] Panasenkov OM, Osipov AN, Chekanov AV, Arnhold J, Sergienko VI. Peroxyl radical is pro-

Diminution of intracellular ROS level by GHK

- duced upon the interaction of hypochlorite with tert-butyl hydroperoxide. *Biochemistry (Moscow)* 2002; 67: 1061-1070.
- [22] Panasenko OM, Chekanov AV, Arnhold J, Sergienko VI, Osipov AN, Vladimirov YA. Generation of free radicals during decomposition of hydroperoxide in the presence of myeloperoxidase or activated neutrophils. *Biochemistry (Moscow)* 2005; 70: 998-1004.
- [23] Barclay LR, Artz JD, Mowat JJ. Partitioning and antioxidant action of the water-soluble antioxidant, Trolox, between the aqueous and lipid phases of phosphatidylcholine membranes: ¹⁴C tracer and product studies. *Biochim Biophys Acta* 1995; 1237: 77-85.
- [24] Stefek M, Kyselova Z, Rackova L, Krizanova L. Oxidative modification of rat eye lens proteins by peroxy radicals in vitro: protection by the chain-breaking antioxidants stobadine and Trolox. *Biochim Biophys Acta* 2005; 1741: 183-190.
- [25] Pickart L, Thaler MM, Millard M. Effect of transition metals on recovery from plasma of the growth-modulating tripeptide glycylhistidyllysine. *J Chromatogr* 1979; 175: 65-73.